

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
29 April 2004 (29.04.2004)

PCT

(10) International Publication Number  
**WO 2004/035047 A1**


- (51) International Patent Classification<sup>7</sup>: **A61K 31/404**, C07D 209/08, 209/30, A61P 25/00 (74) Agents: WAECHTER, Dieter et al.; Grenzacherstrasse 124, CH-4070 Basel (CH).
- (21) International Application Number: PCT/EP2003/011323
- (22) International Filing Date: 13 October 2003 (13.10.2003)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data: 60/419,369 18 October 2002 (18.10.2002) US
- (71) Applicant: F. HOFFMANN-LA ROCHE AG [CH/CH]; 124 Grenzacherstrasse, CH-4070 Basle (CH).
- (72) Inventors: CLARK, Robin, Douglas; 4894 Kua Road, Lawai, HI 96765 (US). HARRIS, Ralph, New, III; 65-D Clinton Street, Redwood City, CA 94062 (US). REPKE, David, Bruce; 406 South Temple Drive, Milpitas, CA 95035 (US).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

**Published:**

— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: 4-PIPERAZINYL BENZENESULFONYL INDOLES WITH 5-HT<sub>6</sub> RECEPTOR AFFINITY



(f)

(57) Abstract: This invention relates to compounds which have generally 5-HT<sub>6</sub> receptor affinity and which are represented by Formula (f): wherein R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup>, and R<sup>5</sup> are as defined herein; or individual isomer, racemic or non-racemic mixtures of isomers, or pharmaceutically acceptable salts or solvates thereof. The invention further relates to pharmaceutical compositions containing such compounds, methods for their use as therapeutic agents, and methods of preparation thereof.


WO 2004/035047 A1

4-PIPERAZINYL BENZENESULFONYL INDOLES WITH 5-HT<sub>6</sub> RECEPTOR AFFINITY

This invention relates to new 4-piperazinyl indole derivatives with 5-HT<sub>6</sub> receptor affinity, and associated pharmaceutical compositions, methods for use as therapeutic agents, and methods of preparation thereof.

The actions of the neurotransmitter 5-hydroxytryptamine (5-HT) as a major modulatory neurotransmitter in the brain, are mediated through a number of receptor families termed 5-HT<sub>1</sub>, 5-HT<sub>2</sub>, 5-HT<sub>3</sub>, 5-HT<sub>4</sub>, 5-HT<sub>5</sub>, 5-HT<sub>6</sub>, and 5-HT<sub>7</sub>. Based on a high level of 5-HT<sub>6</sub> receptor mRNA in the brain, it has been stated that the 5-HT<sub>6</sub> receptor may play a role in the pathology and treatment of central nervous system disorders. In particular, 5-HT<sub>6</sub> receptor selective ligands have been identified as potentially useful in the treatment of certain CNS disorders such as Parkinson's disease, Huntington's disease, anxiety, depression, manic depression, psychoses, epilepsy, obsessive compulsive disorders, migraine, Alzheimer's disease (enhancement of cognitive memory), sleep disorders, feeding disorders such as anorexia and bulimia, panic attacks, attention deficit hyperactivity disorder (ADHD), attention deficit disorder (ADD), withdrawal from drug abuse such as cocaine, ethanol, nicotine, and benzodiazepines, schizophrenia, and also disorders associated with spinal trauma and/or head injury such as hydrocephalus. Such compounds are also expected to be of use in the treatment of certain gastrointestinal (GI) disorders such as functional bowel disorder. (See for ex. B.L. Roth et al., *J. Pharmacol. Exp. Ther.*, 268, pages 1403-14120 (1994), D. R. Sibley et al., *Mol. Pharmacol.*, 43, 320-327 (1993), A.J. Sleight et al, *Neurotransmission*, 11, 1-5 (1995), and A. J. Sleight et al. Serotonin ID Research Alert, 1997, 2 (3), 115-8). 5-HT<sub>6</sub> antagonists have also been identified as potentially useful compounds for treatment of obesity. See for example, Bentley et al., *Br. J. Pharmac.* 1999, *Suppl* 126; Bently et al., *J. Psychopharmacol.* 1997, *Suppl* A64: 255; Wooley et al., *Neuropharmacology* 2001, 41: 210-129; and WO 02/098878.

One object of the present invention is (i) A compound of the formula:



wherein:

$R^1$  and  $R^2$  each independently is hydrogen, alkyl, aryl, acyl, halo, nitro, amino, cyano, alkoxy, hydroxy, aryloxy, alkylthio, arylthio, thiol, carbonylamino, aminocarbonyl, or haloalkyl;

$R^3$  and  $R^4$  each independently is hydrogen, halo, alkyl, acyl, aryl, or arylalkyl;

$n$  is 0 to 4;

$R^6$  in each independent occurrence is hydrogen, alkyl, alkoxy, or halo;

$R^7$  and  $R^8$  each independently is hydrogen or alkyl; and

$R^9$  is hydrogen, alkyl, or arylalkyl;

or an individual isomer, a racemic or non racemic mixture of isomers, a prodrug, or a pharmaceutically acceptable salt or solvate thereof.

Further objects of the present invention are:

(ii) The compound of formula I of (i), wherein:

$R^1$  and  $R^2$  each independently is hydrogen, halo, or alkoxy;


$R^3$  and  $R^4$  each independently is hydrogen, halo or alkyl;

$n$  is 1 and  $R^6$  is hydrogen or halo;

$R^7$  and  $R^8$  are hydrogen; and


$R^9$  is hydrogen or alkyl.

(iii) The compound of formula I of (i), wherein said compound has the formula:




wherein  $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^4$ ,  $n$ ,  $R^6$ ,  $R^7$ ,  $R^8$ , and  $R^9$  are defined in (i).

(iv) The compound of formula I of (i), wherein said compound has the formula:



5 wherein  $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^4$ ,  $n$ ,  $R^6$ ,  $R^7$ ,  $R^8$ , and  $R^9$  are defined in (i).

(v) The compound of formula I of (i), wherein said compound has the formula:



wherein  $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^4$ ,  $n$ ,  $R^6$ ,  $R^7$ ,  $R^8$ , and  $R^9$  are defined in (i).

(vi) The compound of formula Ia of (iii), wherein  $R^1$  and  $R^2$  each independently is  
 10 hydrogen, halo or alkoxy,  $R^3$  is hydrogen or halo;  $R^4$  is hydrogen or alkyl;  $R^6$ ,  $R^7$  and  $R^8$  are hydrogen and  $R^9$  is hydrogen or alkyl.

(vii) The compound of formula Ia of (vi), wherein  $R^1$  and  $R^2$  each independently is hydrogen, fluoro, bromo or methoxy;  $R^3$  is hydrogen or chloro;  $R^4$  each is hydrogen or methyl;  $R^6$ ,  $R^7$  and  $R^8$  are hydrogen and  $R^9$  is hydrogen or methyl.

15 (viii) The compound of formula Ia of (vii), which compound is:

- 5-Fluoro-1-(4-piperazin-1-yl-benzenesulfonyl)-1*H*-indole;  
4-bromo-1-(4-piperazin-1-yl-benzenesulfonyl)-1*H*-indole;  
5-methoxy-1-(4-piperazin-1-yl-benzenesulfonyl)-1*H*-indole;  
1-(4-piperazin-1-yl-benzenesulfonyl)-1*H*-indole;  
5-fluoro-1-[4-(4-methyl-piperazin-1-yl)-benzenesulfonyl]-1*H*-indole;  
3-chloro-5-methoxy-2-methyl-1-(4-piperazin-1-yl-benzenesulfonyl)-1*H*-indole.

(ix) The compound of formula Ib of (iv), wherein  $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^4$ ,  $R^6$ ,  $R^7$ ,  $R^8$  and  $R^9$  are hydrogen.

(x) The compound of formula Ib of (ix), which compound is:

- 2-(4-Piperazin-1-yl-benzenesulfonyl)-1*H*-indole.

(xi) The compound of formula Ic of (v), wherein  $R^1$  and  $R^2$  each independently is hydrogen, halo or alkoxy;  $R^3$  is hydrogen;  $R^4$  is hydrogen or alkyl;  $n$  is 1 and  $R^6$  is hydrogen or halo;  $R^7$  and  $R^8$  are hydrogen and  $R^9$  is hydrogen or alkyl.


- (xii) The compound of formula Ic of (xi), wherein  $R^1$  and  $R^2$  each independently is hydrogen, fluoro, chloro, bromo, iodo or methoxy;  $R^3$  is hydrogen;  $R^4$  is hydrogen or methyl;  $n$  is 1 and  $R^6$  is hydrogen or chloro;  $R^7$  and  $R^8$  are hydrogen and  $R^9$  is hydrogen or methyl.

(xiii) The compound of formula Ic of (xii), which compound is:

- 5-Bromo-1-methyl-3-(4-piperazin-1-yl-benzenesulfonyl)-1*H*-indole;  
5-bromo-3-(4-piperazin-1-yl-benzenesulfonyl)-1*H*-indole;  
3-(4-piperazin-1-yl-benzenesulfonyl)-1*H*-indole;  
1-methyl-3-(4-piperazin-1-yl-benzenesulfonyl)-1*H*-indole;  
6-chloro-3-(4-piperazin-1-yl-benzenesulfonyl)-1*H*-indole;  
4-chloro-3-(4-piperazin-1-yl-benzenesulfonyl)-1*H*-indole;  
5-methoxy-3-(4-piperazin-1-yl-benzenesulfonyl)-1*H*-indole;  
5-chloro-3-(4-piperazin-1-yl-benzenesulfonyl)-1*H*-indole;  
7-chloro-3-(4-piperazin-1-yl-benzenesulfonyl)-1*H*-indole;  
6-chloro-1-methyl-3-(4-piperazin-1-yl-benzenesulfonyl)-1*H*-indole;  
3-[4-(4-methyl-piperazin-1-yl)-benzenesulfonyl]-1*H*-indole;  
7-bromo-3-[4-(4-methyl-piperazin-1-yl)-benzenesulfonyl]-1*H*-indole;  
7-bromo-3-(4-piperazin-1-yl-benzenesulfonyl)-1*H*-indole;  
7-bromo-3-(4-piperazin-1-yl-benzenesulfonyl)-1*H*-indole;  
6-bromo-3-[4-(4-methyl-piperazin-1-yl)-benzenesulfonyl]-1*H*-indole;  
5-iodo-3-[4-(4-methyl-piperazin-1-yl)-benzenesulfonyl]-1*H*-indole;

- 4-bromo-3-[4-(4-methyl-piperazin-1-yl)-benzenesulfonyl]-1H-indole;  
 4-bromo-3-(4-piperazin-1-yl-benzenesulfonyl)-1H-indole;  
 5-iodo-3-(4-piperazin-1-yl-benzenesulfonyl)-1H-indole;  
 6-bromo-3-(4-piperazin-1-yl-benzenesulfonyl)-1H-indole;  
 5 3-(2-chloro-4-piperazin-1-yl-benzenesulfonyl)-1H-indole;  
 3-(3-chloro-4-piperazin-1-yl-benzenesulfonyl)-1H-indole;  
 5-bromo-3-[2-chloro-4-(4-methyl-piperazin-1-yl)-benzenesulfonyl]-1H-indole;  
 5-bromo-3-(2-chloro-4-piperazin-1-yl-benzenesulfonyl)-1H-indole;  
 5-chloro-3-(3-chloro-4-piperazin-1-yl-benzenesulfonyl)-1H-indole;  
 10 5-chloro-3-(3-chloro-4-(4-methyl-piperazin-1-yl)-benzenesulfonyl)-1H-indole;  
 3-(3-chloro-4-(4-methyl-piperazin-1-yl)-benzenesulfonyl)-1H-indole;  
 5-bromo-3-(3-chloro-4-(4-methyl-piperazin-1-yl)-benzenesulfonyl)-1H-indole;  
 5-bromo-3-(3-chloro-4-piperazin-1-yl-benzenesulfonyl)-1H-indole;  
 5-chloro-3-(2-chloro-4-piperazin-1-yl-benzenesulfonyl)-1H-indole;  
 15 5-chloro-3-(2-chloro-4-(4-methyl-piperazin-1-yl)-benzenesulfonyl)-1H-indole;  
 3-(2-chloro-4-(4-methyl-piperazin-1-yl)-benzenesulfonyl)-1H-indole;  
 5-bromo-3-(2-chloro-4-(4-methyl-piperazin-1-yl)-benzenesulfonyl)-1H-indole.


(xiv) A process for producing a compound of formula I of (i), comprising contacting a 4-halobenzenesulfonyl-indole of the formula:



20


wherein X is halo,

with a piperazine of the formula:




to produce a compound of the formula:

- 6 -




wherein  $n$ ,  $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^4$ ,  $R^6$ ,  $R^7$ ,  $R^8$  and  $R^9$  are as defined in (i).

(xv) A process for producing a compound of formula Ia of (iii), comprising contacting formula




wherein X is halo,

with a piperazine of the formula:




to produce a compound of the formula:




wherein  $n$ ,  $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^4$ ,  $R^6$ ,  $R^7$ ,  $R^8$  and  $R^9$  are as defined in (i).

(xvi) A process for producing a compound of formula Ib of (iv), comprising contacting formula




wherein X is halo,

5 with a piperazine of the formula:




to produce a compound of the formula:



wherein n, R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup>, R<sup>6</sup>, R<sup>7</sup>, R<sup>8</sup> and R<sup>9</sup> are as defined in (i).

10 (xvii) A process for producing a compound of formula Ic of (v), comprising contacting formula




wherein X is halo,


with a piperazine of the formula:



- 8 -



to produce a compound of the formula:



wherein  $n$ ,  $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^4$ ,  $R^6$ ,  $R^7$ ,  $R^8$  and  $R^9$  are as defined in (i).

- 5 (xviii) A pharmaceutical composition comprising a therapeutically effective amount of at least one compound of (i) to (xiii) in admixture with one or more pharmaceutically acceptable carrier for the treatment of diseases.

- (xix) Use of one or more compounds of formula I of (i) for the manufacture of a medicament for the treatment or prevention of a disease state that is alleviated by 5-HT<sub>6</sub> agonist.

(xx) The use of (xix)), wherein the disease state comprises disorders of CNS.

- (xxi) The use of (xx), wherein the disease state comprises psychoses, schizophrenia, manic depressions, neurological disorders, memory disorders, attention deficit disorder, Parkinson's disease, amyotrophic lateral sclerosis, Alzheimer's disease and Huntington's disease.

(xxii) The use of (xix), wherein the disease state comprises disorders of the gastrointestinal tract.

(xxiii) The use of (xix), wherein the disease state comprises obesity.

- Unless otherwise stated, the following terms used in this Application, including the specification and claims, have the definitions given below. It must be noted that, as used in the specification and the appended claims, the singular forms "a", "an," and "the" include plural referents unless the context clearly dictates otherwise.

"Alkyl" means the monovalent linear, branched, cyclic, or a combination of linear, branched, or cyclic saturated hydrocarbon radical, consisting solely of carbon and hydrogen atoms, having from one to twelve carbon atoms inclusive, unless otherwise indicated. Examples of alkyl radicals include, but are not limited to, methyl, ethyl, propyl, isopropyl, isobutyl, sec-butyl, tert-butyl, pentyl, n-hexyl, octyl, dodecyl, and the like.

"Lower alkyl" means the monovalent linear or branched saturated hydrocarbon radical, consisting solely of carbon and hydrogen atoms, having from one to six carbon atoms inclusive, unless otherwise indicated. Examples of lower alkyl radicals include, but are not limited to, methyl, ethyl, propyl, isopropyl, sec-butyl, tert-butyl, n-butyl, n-pentyl, n-hexyl, and the like.

"Alkylene" means the divalent linear or branched saturated hydrocarbon radical, consisting solely of carbon and hydrogen atoms, having from one to six carbons inclusive, unless otherwise indicated. Examples of alkylene radicals include, but are not limited to, methylene, ethylene, propylene, 2-methyl-propylene, butylene, 2-ethylbutylene, and the like.

"Alkoxy" means the radical -O-R, wherein R is a lower alkyl radical as defined herein. Examples of alkoxy radicals include, but are not limited to, methoxy, ethoxy, isopropoxy, and the like.

"Halo", "halide" and "halogen", which may be used interchangeably, means fluoro, chloro, bromo, or iodo.

"Haloalkyl" means alkyl substituted with one or more same or different halo atoms, e.g., -CH<sub>2</sub>Cl<sub>2</sub>, -CF<sub>3</sub>, -CH<sub>2</sub>CF<sub>3</sub>, -CH<sub>2</sub>CCl<sub>3</sub>, and the like.

"Alkylthio" or "alkylsulfanyl" means the radical -S-R, wherein R is a lower alkyl radical as defined herein. Examples of alkylthio radicals include, but are not limited to, methylthio, ethylthio, butylthio, and the like.

"Alkylsulfonyl" means the radical -SO<sub>2</sub>R, wherein R is a lower alkyl radical as defined herein. Examples of alkylsulfonyl radicals include, but are not limited to, methylsulfonyl, ethylsulfonyl, and the like.

"Arylthio" or "arylsulfanyl" means the radical -S-R, wherein R is an aryl radical as defined herein.

"Arylsulfonyl" means the radical  $-SO_2R$ , wherein R is an aryl radical as defined herein.

"Aryl" means the monovalent cyclic aromatic hydrocarbon radical consisting of one or more fused rings in which at least one ring is aromatic in nature, which can optionally be substituted with hydroxy, cyano, lower alkyl, lower alkoxy, thioalkyl, halo, haloalkyl, hydroxyalkyl, nitro, alkoxycarbonyl, amino, alkylamino, dialkylamino, aminocarbonyl, carbonylamino, aminosulfonyl, sulfonylamino, and/or trifluoromethyl, unless otherwise indicated. Examples of aryl radicals include, but are not limited to, phenyl, naphthyl, biphenyl, indanyl, anthraquinolyl, and the like. Examples of substituted aryl radicals include but are not limited to fluorophenyl, chlorophenyl, dichlorophenyl, trifluoromethylphenyl, tolyl, and the like.

"Heteroaryl" means the monovalent aromatic carbocyclic radical having one or more rings incorporating one, two, or three heteroatoms within the ring (chosen from nitrogen, oxygen, or sulfur) which can optionally be substituted with hydroxy, cyano, lower alkyl, lower alkoxy, thioalkyl, halo, haloalkyl, hydroxyalkyl, nitro, alkoxycarbonyl, amino, alkylamino, dialkylamino, aminocarbonyl, carbonylamino, aminosulfonyl, sulfonylamino, and/or trifluoromethyl, unless otherwise indicated. Examples of heteroaryl radicals include, but are not limited to, imidazolyl, oxazolyl, thiazolyl, pyrazinyl, thiophenyl, furanyl, pyranyl, pyridinyl, quinolinyl, isoquinolinyl, benzofuryl, benzothiophenyl, benzothiopyranyl, benzimidazolyl, benzooxazolyl, benzothiazolyl, benzopyranyl, indazolyl, indolyl, isoindolyl, naphthyridinyl, and the like.

"Leaving group" means the group with the meaning conventionally associated with it in synthetic organic chemistry, i.e., an atom or group displaceable under alkylating conditions. Examples of leaving groups include, but are not limited to, halogen, alkyl, or arylsulfonyloxy, such as methanesulfonyloxy, ethanesulfonyloxy, thiomethyl, benzenesulfonyloxy, tosyloxy, and thienyloxy, dihalophosphinoyloxy, optionally substituted benzyloxy, isopropoxy, acyloxy, and the like.

"Amino-protecting group" means the protecting group that refers to those organic groups intended to protect the nitrogen atom against undesirable reactions during synthetic procedures and includes, but is not limited to, benzyl (Bnz), benzyloxycarbonyl (carbobenzyloxy, Cbz), p-methoxybenzyloxycarbonyl, p-nitrobenzyloxycarbonyl, *tert*-butoxycarbonyl (Boc), trifluoroacetyl, and the like. It is preferred to use either Boc or Cbz as the amino-protecting group because of the relative ease of removal, for example

by mild acids in the case of Boc, *e.g.*, trifluoroacetic acid or hydrochloric acid in ethyl acetate; or by catalytic hydrogenation in the case of Cbz.

"Optional" or "optionally" means that the subsequently described event or circumstance may but need not occur, and that the description includes instances where the event or circumstance occurs and instances in which it does not. For example, "optional bond" means that the bond may or may not be present, and that the description includes single, double, or triple bonds.

"Protective group" or "protecting group" means the group which selectively blocks one reactive site in a multifunctional compound such that a chemical reaction can be carried out selectively at another unprotected reactive site in the meaning conventionally associated with it in synthetic chemistry. Certain processes of this invention rely upon protective groups to block reactive oxygen atoms present in the reactants. Acceptable protective groups for alcoholic or phenolic hydroxyl groups, which may be removed successively and selectively includes groups protected as acetates, haloalkyl carbonates, benzyl ethers, alkylsilyl ethers, heterocyclyl ethers, methyl or alkyl ethers, and the like. Protective or blocking groups for carboxyl groups are similar to those described for hydroxyl groups, preferably *tert*-butyl, benzyl, or methyl esters.

"Inert organic solvent" or "inert solvent" means the solvent is inert under the conditions of the reaction being described in conjunction therewith, including for example, benzene, toluene, acetonitrile (ACN), tetrahydrofuran (THF), N,N-dimethylformamide (DMF), chloroform, methylene chloride or dichloromethane (DCM), dichloroethane (DCE), diethyl ether, ethyl acetate, acetone, methyl ethyl ketone, methanol, ethanol, propanol, isopropanol, *tert*-butanol, dioxane, pyridine, and the like. Unless specified to the contrary, the solvents used in the reactions of the present invention are inert solvents.

"Pharmaceutically acceptable" means that which is useful in preparing a pharmaceutical composition that is generally safe, non-toxic, and neither biologically nor otherwise undesirable and includes that which is acceptable for veterinary as well as human pharmaceutical use.

"Pharmaceutically acceptable salts" of a compound means salts that are pharmaceutically acceptable, as defined herein, and that possess the desired pharmacological activity of the parent compound. Such salts include:

- (1) acid addition salts formed with inorganic acids such as hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid, and the like; or formed with organic acids such as acetic acid, benzenesulfonic acid, benzoic acid, camphorsulfonic acid, citric acid, ethanesulfonic acid, fumaric acid, glucoheptonic acid, gluconic acid, glutamic acid, glycolic acid, hydroxynaphtoic acid, 2-hydroxyethanesulfonic acid, lactic acid, maleic acid, malic acid, malonic acid, mandelic acid, methanesulfonic acid, muconic acid, 2-naphthalenesulfonic acid, propionic acid, salicylic acid, succinic acid, tartaric acid, p-toluenesulfonic acid, trimethylacetic acid, trifluoroacetic acid, and the like; or
- (2) salts formed when an acidic proton present in the parent compound either is replaced by a metal ion, e.g., an alkali metal ion, an alkaline earth ion, or an aluminum ion; or coordinates with an organic or inorganic base. Acceptable organic bases include diethanolamine, ethanolamine, N-methylglucamine, triethanolamine, tromethamine, and the like. Acceptable inorganic bases include aluminum hydroxide, calcium hydroxide, potassium hydroxide, sodium carbonate, and sodium hydroxide.

The preferred pharmaceutically acceptable salts are the salts formed from acetic acid, trifluoroacetic acid, hydrochloric acid, sulphuric acid, methanesulfonic acid, maleic acid, phosphoric acid, tartaric acid, citric acid, sodium, potassium, calcium, zinc, and magnesium.

- It should be understood that all references to pharmaceutically acceptable salts include solvent addition forms (solvates) or crystal forms (polymorphs) as defined herein, of the same acid addition salt.

- "Solvates" means solvent addition forms that contain either stoichiometric or non stoichiometric amounts of solvent. Some compounds have a tendency to trap a fixed molar ratio of solvent molecules in the crystalline solid state, thus forming a solvate. If the solvent is water the solvate formed is a hydrate, when the solvent is alcohol, the solvate formed is an alcoholate. Hydrates are formed by the combination of one or more molecules of water with one of the substances in which the water retains its molecular state as H<sub>2</sub>O, such combination being able to form one or more hydrate.

- "Prodrug" or "pro-drug" means a pharmacologically inactive form of a compound which must be metabolized in vivo, e.g., by biological fluids or enzymes, by a subject after administration into a pharmacologically active form of the compound in order to produce the desired pharmacological effect. Prodrugs of a compound of Formula I are

prepared by modifying one or more functional group(s) present in the compound of Formula I in such a way that the modification(s) may be cleaved in vivo to release the parent compound. Prodrugs include compounds of Formula I wherein a hydroxy, amino, sulfhydryl, carboxy, or carbonyl group in a compound of Formula I is bonded to  
5 any group that may be cleaved in vivo to regenerate the free hydroxyl, amino, sulfhydryl, carboxy, or carbonyl group respectively. Examples of prodrugs include, but are not limited to, esters (e.g. acetate, dialkylaminoacetates, formates, phosphates, sulfates, and benzoate derivatives), carbamates of hydroxy functional groups ( e.g. N,N-dimethylcarbonyl), esters of carboxyl functional groups (e.g. ethyl esters and  
10 morpholinoethanol esters), N-acyl derivatives (e.g. N-acetyl), N-Mannich bases, Schiff bases, enamines of amino functional groups, oximes, acetals, ketals, enol esters of ketones, aldehyde functional groups in compounds of Formula I, and the like.

The prodrug can be metabolized before absorption, during absorption, after absorption, or at a specific site. Although metabolism occurs for many compounds  
15 primarily in the liver, almost all other tissues and organs, especially the lung, are able to carry out varying degrees of metabolism. Prodrug forms of compounds may be utilized, for example, to improve bioavailability, improve subject acceptability such as by masking or reducing unpleasant characteristics such as bitter taste or gastrointestinal irritability, alter solubility such as for intravenous use, provide for prolonged or sustained release or  
20 delivery, improve ease of formulation, or provide site-specific delivery of the compound. Reference to a compound herein includes prodrug forms of a compound. Prodrugs are described in *The Organic Chemistry of Drug Design and Drug Action*, by Richard B. Silverman, Academic Press, San Diego, 1992. Chapter 8: "Prodrugs and Drug delivery Systems" pp.352-401; *Design of Prodrugs*, edited by H. Bundgaard, Elsevier Science,  
25 Amsterdam, 1985; *Design of Biopharmaceutical Properties through Prodrugs and Analogs*, Ed. by E. B. Roche, American Pharmaceutical Association, Washington, 1977; and *Drug Delivery Systems*, ed. by R.L. Juliano, Oxford Univ. Press, Oxford, 1980.

"Subject" means mammals and non-mammals. Mammals means any member of the Mammalia class including, but not limited to, humans; non-human primates such as  
30 chimpanzees and other apes and monkey species; farm animals such as cattle, horses, sheep, goats, and swine; domestic animals such as rabbits, dogs, and cats; laboratory animals including rodents, such as rats, mice, and guinea pigs; and the like. Examples of non-mammals include, but are not limited to, birds, and the like. The term "subject" does not denote a particular age or sex.

- "Therapeutically effective amount" means an amount of a compound that, when administered to a subject for treating a disease state, is sufficient to effect such treatment for the disease state. The "therapeutically effective amount" will vary depending on the compound, disease state being treated, the severity or the disease treated, the age and relative health of the subject, the route and form of administration, the judgement of the attending medical or veterinary practitioner, and other factors.


"Disease state" means any disease, condition, symptom, or indication.

Throughout the application the following abbreviations may be used with the following meaning:

10	Alk	Alkyl group
	Boc	<i>N</i> - <i>tert</i> -butoxycarbonyl
	<i>m</i> -CPBA	<i>m</i> -Chloroperbenzoic acid
	DCM	Dichloromethane
	DTB	Di- <i>tert</i> -butyldicarbonate
15	DMF	<i>N,N</i> -Dimethylformamide
	DMFDMA	<i>N,N</i> -dimethylformamide dimethyl acetal
	DMSO	Dimethylsulfoxide
	L	Leaving group
	Oxone <sup>TM</sup>	Potassium peroxymonosulfate
20	P	Protective group
	TFA	Trifluoroacetic acid
	THF	Tetrahydrofuran


The naming and numbering of the compounds of this invention is illustrated below:

- 15 -




In general, the nomenclature used herein is based on AUTONOM<sup>TM</sup> v.4.0, a Beilstein Institute computerized system for the generation of IUPAC systematic nomenclature. However, because a strict adherence to these recommendations would result in the names changing substantially when only a single substituent is changed, compounds have been named in a manner that maintains consistency of nomenclature for the basic structure of the molecule.

For example, a compound of Formula I wherein R<sup>2</sup>, R<sup>3</sup> and R<sup>4</sup> are hydrogen, R<sup>1</sup> is fluorine (attached through the 5<sup>th</sup> position on the indole ring), and the group



is attached through the 1<sup>st</sup> position on the indole ring, R<sup>6</sup>, R<sup>7</sup> and R<sup>8</sup> are hydrogen, and R<sup>9</sup> is methyl, is named 5-Fluoro-1-[4-(4-methyl-piperazin-1-yl)-benzenesulfonyl]-1H-indole.

The invention provides Compounds of Formula I:



wherein:

R<sup>1</sup> and R<sup>2</sup> each independently is hydrogen, alkyl, aryl, acyl, halo, nitro, amino, cyano, alkoxy, hydroxy, aryloxy, alkylthio, arylthio, thiol, carbonylamino,



- 16 -

aminocarbonyl, or haloalkyl; preferably  $R^1$  and  $R^2$  each independently is hydrogen, halo or alkoxy; more preferably hydrogen or halo;

$R^3$  and  $R^4$  each independently is hydrogen, halo, alkyl, acyl, aryl, or arylalkyl; preferably  $R^3$  and  $R^4$  each independently is hydrogen or alkyl;

5  $n$  is from 0 to 3; preferably  $n$  is 0 or 1;


$R^5$  in each occurrence independently is hydrogen, alkyl, alkoxy, or halo; preferably  $R^6$  is hydrogen or halo;

$R^7$  and  $R^8$  each independently is hydrogen or alkyl; and

$R^9$  is hydrogen, alkyl, or arylalkyl; preferably  $R^9$  is hydrogen or alkyl;


10 or individual isomers, racemic or non racemic mixtures of isomers, prodrugs, or pharmaceutically acceptable salts or solvates thereof.

In certain embodiments of the invention, the Compounds of Formula I may be of Formula Ia:




15 wherein  $n$ ,  $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^4$ ,  $R^6$ ,  $R^7$ ,  $R^8$  and  $R^9$  are as defined herein.

In other embodiments, Compounds of Formula I may be of Formula Ib:



wherein  $n$ ,  $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^4$ ,  $R^6$ ,  $R^7$ ,  $R^8$  and  $R^9$  are as defined herein.

In more preferred embodiments, Compounds of Formula I are of the Formula Ic:



Ic








wherein n, R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup>, R<sup>6</sup>, R<sup>7</sup>, R<sup>8</sup> and R<sup>9</sup> are as defined herein.







Representative compounds in accordance with the invention are shown in Table 1.








5 In many instances the compounds in Table 1 are shown as hydrochloride salts.







TABLE 1







#	STRUCTURE	NAME	M+H or MP °C	Example
1		5-Fluoro-1-(4-piperazin-1-yl-benzenesulfonyl)-1H-indole	M+H 360	1
2		4-Bromo-1-(4-piperazin-1-yl-benzenesulfonyl)-1H-indole	M+H 421	1
3		5-Methoxy-1-(4-piperazin-1-yl-benzenesulfonyl)-1H-indole	M+H 372	1
4		1-(4-Piperazin-1-yl-benzenesulfonyl)-1H-indole	M+H 342	1




#	STRUCTURE	NAME	M+H or MP °C	Example
5		5-Fluoro-1-[4-(4-methyl-piperazin-1-yl)-benzenesulfonyl]-1H-indole	M+H 374	1
6		5-Bromo-1-methyl-3-(4-piperazin-1-yl-benzenesulfonyl)-1H-indole	M+H 435	3
7		5-Bromo-3-(4-piperazin-1-yl-benzenesulfonyl)-1H-indole	M+H 421	3
8		3-(4-Piperazin-1-yl-benzenesulfonyl)-1H-indole	M+H 342	3
9		3-Chloro-5-methoxy-2-methyl-1-(4-piperazin-1-yl-benzenesulfonyl)-1H-indole	M+H 420	1
10		1-Methyl-3-(4-piperazin-1-yl-benzenesulfonyl)-1H-indole	M+H 356	3
11		6-Chloro-3-(4-piperazin-1-yl-benzenesulfonyl)-1H-indole	M+H 376	3

#	STRUCTURE	NAME	M+H or MP °C	Example
12		4-Chloro-3-(4-piperazin-1-yl-benzenesulfonyl)-1H-indole	M+H 376	3
13		5-Methoxy-3-(4-piperazin-1-yl-benzenesulfonyl)-1H-indole	M+H 372	3
14		5-Chloro-3-(4-piperazin-1-yl-benzenesulfonyl)-1H-indole	M+H 376	3
15		7-Chloro-3-(4-piperazin-1-yl-benzenesulfonyl)-1H-indole	M+H 376	3
16		6-Chloro-1-methyl-3-(4-piperazin-1-yl-benzenesulfonyl)-1H-indole	M+H 422	3
17		3-[(4-(4-Methylpiperazin-1-yl)-benzenesulfonyl)-1H-indole	M+H 356	3

#	STRUCTURE	NAME	M+H or MP °C	Example
18	 ClH	7-Bromo-3-[4-(4-methyl-piperazin-1-yl)-benzenesulfonyl]-1H-indole	M+H 435	3
19	 ClH	7-Bromo-3-(4-piperazin-1-yl-benzenesulfonyl)-1H-indole	M+H 421	3
20	 ClH	6-Bromo-3-[4-(4-methyl-piperazin-1-yl)-benzenesulfonyl]-1H-indole	M+H 435	3
21	 ClH	5-Iodo-3-[4-(4-methyl-piperazin-1-yl)-benzenesulfonyl]-1H-indole	M+H 482	3
22	 ClH	4-Bromo-3-[4-(4-methyl-piperazin-1-yl)-benzenesulfonyl]-1H-indole	M+H 435	3
23	 ClH	4-Bromo-3-(4-piperazin-1-yl-benzenesulfonyl)-1H-indole	M+H 421	3
24	 ClH	5-Iodo-3-(4-piperazin-1-yl-benzenesulfonyl)-1H-indole	M+H 468	3

#	STRUCTURE	NAME	M+H or MP °C	Example
25		6-Bromo-3-(4-piperazin-1-yl-benzenesulfonyl)-1H-indole	M+H 421	3
26		3-(2-Chloro-4-piperazin-1-yl-benzenesulfonyl)-1H-indole	M+H 376	3
27		3-(3-Chloro-4-piperazin-1-yl-benzenesulfonyl)-1H-indole	M+H 376	3
28		5-Bromo-3-[2-chloro-4-(4-methyl-piperazin-1-yl)-benzenesulfonyl]-1H-indole	M+H 468	3
29		5-Bromo-3-(2-chloro-4-piperazin-1-yl-benzenesulfonyl)-1H-indole	MP 283- 286°	3
30		5-Chloro-3-(3-chloro-4-piperazin-1-yl-benzenesulfonyl)-1H-indole	MP 252-258°	3

#	STRUCTURE	NAME	M+H or MP °C	Example
31		5-Chloro-3-(3-chloro-4-(4-methyl-piperazin-1-yl)-benzenesulfonyl)-1H-indole	MP 198-202°	3
32		3-(3-Chloro-4-(4-methyl-piperazin-1-yl)-benzenesulfonyl)-1H-indole	M+H 390	3
33		5-Bromo-3-(3-chloro-4-(4-methyl-piperazin-1-yl)-benzenesulfonyl)-1H-indole	MP 194-197°	3
34		5-Bromo-3-(3-chloro-4-piperazin-1-yl-benzenesulfonyl)-1H-indole	MP 210-218°	3
35		5-Chloro-3-(2-chloro-4-piperazin-1-yl-benzenesulfonyl)-1H-indole	M+H 410	3
36		5-Chloro-3-(2-chloro-4-(4-methyl-piperazin-1-yl)-benzenesulfonyl)-1H-indole	MP 250°	3

#	STRUCTURE	NAME	M+H or MP °C	Example
37		3-(2-Chloro-4-(4-methyl-piperazin-1-yl)-benzenesulfonyl)-1H-indole	M+H 390	3
38		5-Bromo-3-(2-chloro-4-(4-methyl-piperazin-1-yl)-benzenesulfonyl)-1H-indole	M+H 468	3
39		2-(4-Piperazin-1-yl-benzenesulfonyl)-1H-indole	M+H 342	2

The invention provides, in another aspect, pharmaceutical compositions comprising at least one Compound of Formula I together with one or more pharmaceutically acceptable carriers, diluents, adjuvants or excipients.

- 5 Yet another aspect of the invention provides methods for treating a central nervous system (CNS) disease state or condition in a subject comprising administering to a subject in need thereof a therapeutically effective amount of at least one Compound of Formula I. The disease state may comprise psychoses, schizophrenia, manic depressions, neurological disorders, memory disorders, attention deficit disorder, Parkinson's disease,
- 10 amyotrophic lateral sclerosis, Alzheimer's disease and/or Huntington's disease.

Still another aspect of the present invention provides a method for treating a disorder of the gastrointestinal tract in a subject comprising administering to the subject a therapeutically effective amount of a Compound of Formula I.

- Another aspect of the present invention provides methods for producing
- 15 Compounds of Formula I according to the synthetic procedures described below.

Compounds of the present invention may be made by the methods depicted in the illustrative synthetic reaction schemes shown and described below.




The starting materials and reagents used in preparing these compounds generally are either available from commercial suppliers, such as Aldrich Chemical Co., or are prepared by methods known to those skilled in the art following procedures set forth in references such as *Fieser and Fieser's Reagents for Organic Synthesis*, Wiley & Sons: New York, 1991, Volumes 1-15; *Rodd's Chemistry of Carbon Compounds*, Elsevier Science Publishers, 1989, Volumes 1-5 and Supplementals; and *Organic Reactions*, Wiley & Sons: New York, 1991, Volumes 1-40. The indole starting materials used herein are commercially available or may be prepared by conventional techniques such as those described by Sundberg, R.J. in *The Chemistry of Indoles*, Academic Press, New York, 1970.

The following synthetic reaction schemes are merely illustrative of some methods by which the compounds of the present invention may be synthesized, and various modifications to these synthetic reaction schemes may be made and will be suggested to one skilled in the art having referred to the disclosure contained in this Application.

The starting materials and the intermediates of the synthetic reaction schemes may be isolated and purified if desired using conventional techniques, including but not limited to filtration, distillation, crystallization, chromatography, and the like. Such materials may be characterized using conventional means, including physical constants and spectral data.

Unless specified to the contrary, the reactions described herein preferably take place at atmospheric pressure over a temperature range from about -78° C to about 150° C, more preferably from about 0° C to about 125° C, and most preferably and conveniently at about room (or ambient) temperature, e.g., about 20° C.


Compounds of Formula Ia may be prepared as shown in Scheme A wherein X, n, R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup>, R<sup>6</sup>, R<sup>7</sup>, R<sup>8</sup> and R<sup>9</sup> are as defined herein.



## SCHEME A

In Scheme A, the 1-(4-halobenzenesulfonyl)-1*H*-indole **c** is formed by reaction of the indole compound **f** under basic conditions with a 4-halophenylsulfonyl halide **g** having a desired R<sup>6</sup> substituent(s). The 1-(4-halobenzenesulfonyl)-1*H*-indole **c** may then be treated with excess piperazine **h** in polar aprotic solvent to afford the corresponding 1-(4-(piperazin-1-yl)-benzenesulfonyl)-1*H*-indole of Formula Ia.


Compounds of Formula Ib may be prepared according to the procedure shown in Scheme B, wherein X, n, R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup>, R<sup>6</sup>, R<sup>7</sup>, R<sup>8</sup> and R<sup>9</sup> are as defined herein.



## SCHEME B

In Scheme B, the indole compound **h** is treated with an alkyllithium reagent or other strong base under anhydrous polar aprotic conditions and dry ice/acetone temperature to generate the corresponding indole anion (not shown) by deprotonation at the 2- position. Where R<sup>4</sup> is hydrogen, a suitable removable protecting group may be used to protect the indole nitrogen of compound **h**. A 4-halophenylsulfonyl fluoride **g** may then be added directly to the anion of indole compound **h** to provide the 2-(4-halobenzenesulfonyl)-1*H*-indole **d**. Treatment of the 2-(4-halobenzenesulfonyl)-1*H*-indole **d** with excess piperazine **b** can then provide the corresponding 2-(4-(piperazin-1-yl)-benzenesulfonyl)-1*H*-indole of Formula Ib.

Compounds of Formula Ic may be prepared according to the procedure shown in Scheme C, wherein X, n, R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup>, R<sup>6</sup>, R<sup>7</sup>, R<sup>8</sup> and R<sup>9</sup> again are as defined herein.



SCHEME C

In Scheme C, the indole compound **i** is reacted with a 4-halophenyl disulfide **j** in the presence of alkalai metal hydride or like strong base under dry polar aprotic conditions to yield the 3-(4-halophenylsulfanyl)-1H-indole compound **k**. In the procedure of Scheme C, R<sup>4</sup> of compound **i** is preferably hydrogen. The compound **k** may then be treated with a peracid or like oxidizing agent to afford the 3-(4-halobenzenesulfonyl)-1H-indole compound **e**. Treatment of the 3-(4-halobenzenesulfonyl)-1H-indole **e** with excess piperazine **b** furnishes the 3-(4-piperazin-1-yl-benzenesulfonyl)-1H-indole of Formula 1c. Where it is desired to have R<sup>4</sup> as alkyl in compounds of Formula 1c, alkylation of the indole nitrogen at the 1- position of the indole ring may be carried out subsequent to formation of the sulfanyl compound **k**.

Many variations of the above synthetic schemes are possible and will suggest themselves to those skilled in the art. Those skilled in the art will also recognize that stereocenters exist in some compounds of Formula I. Accordingly, the present invention includes all possible stereoisomers and geometric isomers of Formula I, and includes not only racemic compounds but also the optically active isomers as well. When a Compound of Formula I, is desired as a single enantiomer, it may be obtained either by resolution of the final product or by stereospecific synthesis from either isomerically pure starting material or any convenient intermediate. Resolution of the final product, an intermediate or a starting material may be effected by any suitable method known in the art. See for example, *Stereochemistry of Carbon Compounds* by E.L.Eliel (McGraw Hill, 1962) and *Tables of Resolving Agents* by S.H.Wilen.

The compounds of the invention have selective 5-HT<sub>6</sub> receptor affinity and as such are expected to be useful in the treatment of certain CNS disorders such as Parkinson's

- disease, Huntington's disease, anxiety, depression, manic depression, psychosis, epilepsy, obsessive compulsive disorders, migraine, Alzheimer's disease (enhancement of cognitive memory), sleep disorders, feeding disorders such as anorexia and bulimia, panic attacks, attention deficit hyperactivity disorder (ADHD), attention deficit disorder (ADD),
- 5 withdrawal from drug abuse such as cocaine, ethanol, nicotine, and benzodiazepines, schizophrenia, disorders associated with spinal trauma and/or head injury such as hydrocephalus, and other diseases, disorders or conditions mediated by or otherwise associated with the 5-HT<sub>6</sub> receptor or other 5-HT receptors. Such compounds are also expected to be of use in the treatment of certain GI (gastrointestinal) disorders such as
- 10 functional bowel disorder or irritable bowel syndrome (IBS).

The pharmacology of the compounds of this invention was determined by art recognized procedures. The in vitro techniques for determining the affinities of test compounds at the 5-HT<sub>6</sub> receptor in radioligand binding and functional assays are described in Example 5.

- 15 The present invention includes pharmaceutical compositions comprising at least one compound of the present invention, or an individual isomer, racemic or non-racemic mixture of isomers or a pharmaceutically acceptable salt or solvate thereof, together with at least one pharmaceutically acceptable carrier, and optionally other therapeutic and/or prophylactic ingredients.

- 20 In general, the compounds of the present invention will be administered in a therapeutically effective amount by any of the accepted modes of administration for agents that serve similar utilities. Suitable dosage ranges are typically 1-500 mg daily, preferably 1-100 mg daily, and most preferably 1-30 mg daily, depending upon numerous factors such as the severity of the disease to be treated, the age and relative
- 25 health of the subject, the potency of the compound used, the route and form of administration, the indication towards which the administration is directed, and the preferences and experience of the medical practitioner involved. One of ordinary skill in the art of treating such diseases will be able, without undue experimentation and in reliance upon personal knowledge and the disclosure of this Application, to ascertain a
- 30 therapeutically effective amount of the compounds of the present invention for a given disease.

In general, compounds of the present invention will be administered as pharmaceutical formulations including those suitable for oral (including buccal and sub-lingual), rectal, nasal, topical, pulmonary, vaginal, or parenteral (including

intramuscular, intraarterial, intrathecal, subcutaneous, and intravenous) administration or in a form suitable for administration by inhalation or insufflation. The preferred manner of administration is generally oral using a convenient daily dosage regimen which can be adjusted according to the degree of affliction.

5       A compound or compounds of the present invention, together with one or more conventional adjuvants, carriers, or diluents, may be placed into the form of pharmaceutical compositions and unit dosages. The pharmaceutical compositions and unit dosage forms may be comprised of conventional ingredients in conventional proportions, with or without additional active compounds or principles, and the unit  
10       dosage forms may contain any suitable effective amount of the active ingredient commensurate with the intended daily dosage range to be employed. The pharmaceutical compositions may be employed as solids, such as tablets or filled capsules, semisolids, powders, sustained release formulations, or liquids such as solutions, suspensions, emulsions, elixirs, or filled capsules for oral use; or in the form of suppositories for rectal  
15       or vaginal administration; or in the form of sterile injectable solutions for parenteral use. Formulations containing about one (1) milligram of active ingredient or, more broadly, about 0.01 to about one hundred (100) milligrams, per tablet, are accordingly suitable representative unit dosage forms.

      The compounds of the present invention may be formulated in a wide variety of  
20       oral administration dosage forms. The pharmaceutical compositions and dosage forms may comprise a compound or compounds of the present invention or pharmaceutically acceptable salts thereof as the active component. The pharmaceutically acceptable carriers may be either solid or liquid. Solid form preparations include powders, tablets, pills, capsules, cachets, suppositories, and dispersible granules. A solid carrier may be one  
25       or more substances which may also act as diluents, flavouring agents, solubilizers, lubricants, suspending agents, binders, preservatives, tablet disintegrating agents, or an encapsulating material. In powders, the carrier generally is a finely divided solid which is a mixture with the finely divided active component. In tablets, the active component generally is mixed with the carrier having the necessary binding capacity in suitable  
30       proportions and compacted in the shape and size desired. The powders and tablets preferably contain from about one (1) to about seventy (70) percent of the active compound. Suitable carriers include but are not limited to magnesium carbonate, magnesium stearate, talc, sugar, lactose, pectin, dextrin, starch, gelatine, tragacanth, methylcellulose, sodium carboxymethylcellulose, a low melting wax, cocoa butter, and  
35       the like. The term "preparation" is intended to include the formulation of the active

compound with encapsulating material as carrier, providing a capsule in which the active component, with or without carriers, is surrounded by a carrier, which is in association with it. Similarly, cachets and lozenges are included. Tablets, powders, capsules, pills, cachets, and lozenges may be as solid forms suitable for oral administration.

- 5 Other forms suitable for oral administration include liquid form preparations including emulsions, syrups, elixirs, aqueous solutions, aqueous suspensions, or solid form preparations which are intended to be converted shortly before use to liquid form preparations. Emulsions may be prepared in solutions, for example, in aqueous propylene glycol solutions or may contain emulsifying agents, for example, such as  
10 lecithin, sorbitan monooleate, or acacia. Aqueous solutions can be prepared by dissolving the active component in water and adding suitable colorants, flavors, stabilizing, and thickening agents. Aqueous suspensions can be prepared by dispersing the finely divided active component in water with viscous material, such as natural or synthetic gums, resins, methylcellulose, sodium carboxymethylcellulose, and other well known  
15 suspending agents. Solid form preparations include solutions, suspensions, and emulsions, and may contain, in addition to the active component, colorants, flavors, stabilizers, buffers, artificial and natural sweeteners, dispersants, thickeners, solubilizing agents, and the like.

- The compounds of the present invention may be formulated for parenteral  
20 administration (e.g., by injection, for example bolus injection or continuous infusion) and may be presented in unit dose form in ampoules, pre-filled syringes, small volume infusion or in multi-dose containers with an added preservative. The compositions may take such forms as suspensions, solutions, or emulsions in oily or aqueous vehicles, for example solutions in aqueous polyethylene glycol. Examples of oily or nonaqueous  
25 carriers, diluents, solvents or vehicles include propylene glycol, polyethylene glycol, vegetable oils (e.g., olive oil), and injectable organic esters (e.g., ethyl oleate), and may contain formulatory agents such as preserving, wetting, emulsifying or suspending, stabilising and/or dispersing agents. Alternatively, the active ingredient may be in powder form, obtained by aseptic isolation of sterile solid or by lyophilisation from solution for  
30 constitution before use with a suitable vehicle, e.g., sterile, pyrogen-free water.

- The compounds of the present invention may be formulated for topical  
administration to the epidermis as ointments, creams or lotions, or as a transdermal patch. Ointments and creams may, for example, be formulated with an aqueous or oily  
base with the addition of suitable thickening and/or gelling agents. Lotions may be  
35 formulated with an aqueous or oily base and will in general also containing one or more

emulsifying agents, stabilising agents, dispersing agents, suspending agents, thickening agents, or colouring agents. Formulations suitable for topical administration in the mouth include lozenges comprising active agents in a flavoured base, usually sucrose and acacia or tragacanth; pastilles comprising the active ingredient in an inert base such as gelatine and glycerine or sucrose and acacia; and mouthwashes comprising the active ingredient in a suitable liquid carrier.

The compounds of the present invention may be formulated for administration as suppositories. A low melting wax, such as a mixture of fatty acid glycerides or cocoa butter is first melted and the active component is dispersed homogeneously, for example, by stirring. The molten homogeneous mixture is then poured into convenient sized molds, allowed to cool, and to solidify.

The compounds of the present invention may be formulated for vaginal administration. Pessaries, tampons, creams, gels, pastes, foams or sprays containing in addition to the active ingredient such carriers as are known in the art to be appropriate.

The compounds of the present invention may be formulated for nasal administration. The solutions or suspensions are applied directly to the nasal cavity by conventional means, for example, with a dropper, pipette or spray. The formulations may be provided in a single or multidose form. In the latter case of a dropper or pipette, this may be achieved by the patient administering an appropriate, predetermined volume of the solution or suspension. In the case of a spray, this may be achieved for example by means of a metering atomising spray pump.

The compounds of the present invention may be formulated for aerosol administration, particularly to the respiratory tract and including intranasal administration. The compound will generally have a small particle size for example of the order of five (5) microns or less. Such a particle size may be obtained by means known in the art, for example by micronization. The active ingredient is provided in a pressurized pack with a suitable propellant such as a chlorofluorocarbon (CFC), for example, dichlorodifluoromethane, trichlorofluoromethane, or dichlorotetrafluoroethane, or carbon dioxide or other suitable gas. The aerosol may conveniently also contain a surfactant such as lecithin. The dose of drug may be controlled by a metered valve. Alternatively the active ingredients may be provided in a form of a dry powder, for example a powder mix of the compound in a suitable powder base such as lactose, starch, starch derivatives such as hydroxypropylmethyl cellulose and polyvinylpyrrolidone (PVP). The powder carrier will form a gel in the nasal cavity. The powder composition

may be presented in unit dose form for example in capsules or cartridges of e.g., gelatine or blister packs from which the powder may be administered by means of an inhaler.

When desired, formulations can be prepared with enteric coatings adapted for sustained or controlled release administration of the active ingredient. For example, the compounds of the present invention can be formulated in transdermal or subcutaneous drug delivery devices. These delivery systems are advantageous when sustained release of the compound is necessary and when patient compliance with a treatment regimen is crucial. Compounds in transdermal delivery systems are frequently attached to a skin-adhesive solid support. The compound of interest can also be combined with a penetration enhancer, e.g., Azone (1-dodecylazacycloheptan-2-one). Sustained release delivery systems are inserted subcutaneously into the subdermal layer by surgery or injection. The subdermal implants encapsulate the compound in a lipid soluble membrane, e.g., silicone rubber, or a biodegradable polymer, e.g., polylactic acid.

The pharmaceutical preparations are preferably in unit dosage forms. In such form, the preparation is subdivided into unit doses containing appropriate quantities of the active component. The unit dosage form can be a packaged preparation, the package containing discrete quantities of preparation, such as packeted tablets, capsules, and powders in vials or ampoules. Also, the unit dosage form can be a capsule, tablet, cachet, or lozenge itself, or it can be the appropriate number of any of these in packaged form.

Other suitable pharmaceutical carriers and their formulations are described in *Remington: The Science and Practice of Pharmacy* 1995, edited by E. W. Martin, Mack Publishing Company, 19<sup>th</sup> edition, Easton, Pennsylvania. Representative pharmaceutical formulations containing a compound of the present invention are described in Example 4.

## EXAMPLES


The following preparations and examples are given to enable those skilled in the art to more clearly understand and to practice the present invention. They should not be considered as limiting the scope of the invention, but merely as being illustrative and representative thereof.

### Example 1

5-Fluoro-1-(4-piperazin-1-yl-benzenesulfonyl)-1H-indole




- 32 -



### Step 1

#### 1-(4-Chloro-benzenesulfonyl)-5-fluoro-1H-indole




- 5           A mixture of 5-fluoroindole (1.0g, 7.4mmole), 4-chlorobenzenesulfonyl chloride (1.6g, 7.6mmole), tetra-*n*-butylammonium hydrogen sulfate (0.05g), 4*N* sodium hydroxide (5mL, 20mmol), and toluene (6mL) was stirred at room temperature for 15h. The mixture was diluted with 10mL water and extracted with 25mL ethyl acetate. The organic phase was washed with 10mL water, 10mL saturated sodium chloride, and then
- 10       dried (anhydrous magnesium sulfate). The solution was concentrated under reduced pressure. The residue was recrystallized from ethyl ether/hexane to provide 1-(4-Chloro-benzenesulfonyl)-5-fluoro-1H-indole (1.74g) m.p. 108-109 °C.

### Step 2

#### 5-Fluoro-1-(4-(piperazin-1-yl)-benzenesulfonyl)-1H-indole

- 33 -



A solution of 1-(4-Chlorobenzenesulfonyl)-5-fluoro-1H-indole (0.210g, 0.7mmole) and anhydrous piperazine (0.3 g, 3.4mmole) in dimethylsulfoxide (3.0mL) was heated at 100 °C in a sealed tube for 16h. The reaction mixture was diluted with  
 5 15mL water and extracted with 20mL ethyl acetate. The organic phase was washed with three 10mL portions of water, 10mL saturated sodium chloride, dried (magnesium sulfate), and concentrated under reduced pressure. The residue was recrystallized from ethyl acetate/hexane to provide the 5-Fluoro-1-(4-piperazin-1-yl-benzenesulfonyl)-1H-indole (0.123 g, m.p. 77-78 °C).


10 Similarly following the procedure described above in Example 1, but replacing 5-fluorindole in Step 1 with other appropriate indoles, the following compounds were prepared:


- 4-bromo-1-(4-piperazin-1-yl-benzenesulfonyl)-1H-indole;
- 5-methoxy-1-(4-piperazin-1-yl-benzenesulfonyl)-1H-indole;
- 15 1-(4-piperazin-1-yl-benzenesulfonyl)-1H-indole;
- 5-fluoro-1-[4-(4-methyl-piperazin-1-yl)-benzenesulfonyl]-1H-indole;
- 3-chloro-5-methoxy-2-methyl-1-(4-piperazin-1-yl-benzenesulfonyl)-1H-indole.

Similarly following the procedure described above in Example 1, but replacing  
 20 piperazine in Step 2 with N-methylpiperazine, the compound 5-Fluoro-1-[4-(4-methyl-piperazin-1-yl)-benzenesulfonyl]-1H-indole was prepared.


### Example 2

#### 2-(4-Piperazin-1-yl-benzenesulfonyl)-1H-indole



Step 12-(4-Fluoro-benzenesulfonyl)-indole-1-carboxylic acid *tert*-butyl ester


- A solution of 1-indole carboxylic acid *tert*-butyl ester (1.88 g, 8.65 mmol) in 100 mL of anhydrous THF was cooled to  $-78^{\circ}\text{C}$  in a dry ice/acetone bath under an argon gas blanket. A fresh commercial (Aldrich) 1.7 M solution of *t*-BuLi in pentane (10.2 mL, 17.3 mmol, 2.0 eq) was slowly added to the stirring mixture via syringe. During the addition, the solution turned from a light amber-red color to a dark, orange-burgundy color. After the addition was complete, the reaction mixture was allowed to stir for 45 minutes at  $-78^{\circ}\text{C}$  and then 4-fluorobenzenesulfonyl fluoride was slowly added to the reaction mixture while carefully maintaining an internal temperature of  $<-65^{\circ}\text{C}$ . The reaction mixture was kept at  $-75$  to  $-78^{\circ}\text{C}$  for a period of 20 minutes after the addition was complete then was allowed to warm to room temperature, where it remained for 2h. The reaction mixture was poured into 500 mL of saturated  $\text{NH}_4\text{Cl}$  solution, diluted with 500 mL of EtOAc and mixed well. The pale, amber-orange organic layer was separated, washed with brine and dried over anhydrous magnesium sulfate. Evaporation of the solvent gave a dark amber oil which was silica-gel chromatographed, gradient eluting with 10% EtOAc/Hexane followed by  $\text{CHCl}_3$  to give 1.40g, 3.73 mmol, 43.1% yield of the 2-(4-Fluoro-benzenesulfonyl)-indole-1-carboxylic acid *tert*-butyl ester as an amber oil which solidified on standing to a pale reddish-tan solid which had a melting point range of  $88-93^{\circ}\text{C}$ .

Step 22-(4-Piperazin-1-yl-benzenesulfonyl)-1H-indole

The Boc-protecting group on 0.530 g, 1.41 mmol of the 2-(4-Fluoro-benzenesulfonyl)-indole-1-carboxylic acid *tert*-butyl ester from Step 1 was removed by stirring at room temperature in 3 mL of neat TFA. The reaction mixture turned a light burgundy color while stirring. The TFA was removed by adding 25 mL of toluene and removing the solvent/TFA as an azeotrope. This was repeated and the oily residue was then transferred to a sealed tube using 5 mL of DMSO. Solid piperazine (0.61 g, 7.06 mmol, 5 eq.) was added, the tube was sealed, and the reaction mixture heated at 100°C (bath temp.) for a period of 16 h. The reaction mixture was allowed to cool to room temperature, diluted with 400 mL of ethyl acetate, then washed sequentially with saturated NaHCO<sub>3</sub> solution, water, brine, and dried over anhydrous magnesium sulfate. Evaporation of the solvent gave an amber glass which was treated with anhydrous diethyl ether. Hexane (25 mL) was added to the solution, causing a small amount of solid to triturate out of solution. The mixture cooled was to 0°C in an ice-water bath, followed by vacuum-filtration of the off-white solid. After drying, 0.150g, 0.44 mmol, 31.1% yield of 2-(4-Piperazin-1-yl-benzenesulfonyl)-1H-indole was obtained, with a melting point range of 246-248.5°C and M+H of 342.


### Example 3

#### 5-Bromo-3-(4-piperazin-1-yl-benzenesulfonyl)-1H-indole



### Step 1

#### 5-Bromo-3-(4-fluoro-phenylsulfanyl)-1H-indole




To a stirred mixture of sodium hydride (100%, 0.156g, 6.5mmole) in dry N,N-dimethylformamide (12mL) was added 5-bromoindole (0.98g, 5 mmole). After 5 minutes 4-fluorophenyl disulfide (1.4g, 5.5mmole) was added. The mixture was stirred at room temperature for 48h. The mixture was diluted with 50mL water and extracted with 25mL ethyl ether. The organic phase was washed with 1.5M sodium carbonate (5mL,

- 36 -

7.5mmol), 5mL water, dried (magnesium sulfate), and concentrated under reduced pressure to yield 5-Bromo-3-(4-fluoro-phenylsulfanyl)-1H-indole (1.34g).

### Step 2

#### 5-Bromo-3-(4-fluoro-benzenesulfonyl)-1H-indole




5

The 5-Bromo-3-(4-fluoro-phenylsulfanyl)-1H-indole (1.34g, 4.16mmol) from Step 1 was suspended in formic acid (20mL). To this was added 30% hydrogen peroxide (0.94g, 8.3mmole). The reaction mixture was stirred at room temperature for 1 hour. The mixture was diluted with 80mL water and extracted with 100mL ethyl acetate. The organic phase was washed with 1.5M sodium carbonate (20mL, 30mmol), dried (magnesium sulfate), and concentrated under reduced pressure. The residue was recrystallized from ether/hexane to provide 5-Bromo-3-(4-fluoro-benzenesulfonyl)-1H-indole (1.4 g, 4.0mmol), m.p. 163-164 °C.

### Step 3

#### 5-Bromo-3-(4-fluoro-benzenesulfonyl)-1-methyl-1H-indole



15


To a stirred solution of 5-bromo-3-(4-fluoro-benzenesulfonyl)-1H-indole (0.315g, 0.89mmole) in tetrahydrofuran (4mL) was added 1M potassium t-butoxide/THF (1.0mL, 1.0mmol). After 10 minutes iodomethane (0.16g, 1.1mmole) was added. The reaction mixture was stirred at room temperature overnight and then passed through a pad of silica gel (230-400 mesh) eluting with 50% ethyl acetate/hexane. The eluate was concentrated under reduced pressure to provide 5-Bromo-3-(4-fluoro-benzenesulfonyl)-1-methyl-1H-indole.

20

### Step 4

#### 5-Bromo-1-methyl-3-(4-piperazin-1-yl-benzenesulfonyl)-1H-indole

25



The 5-Bromo-3-(4-fluoro-benzenesulfonyl)-1-methyl-1H-indole (0.30 g, 0.85 mmol) was mixed with piperazine (0.34g, 4mmole) and methyl sulfoxide (3mL). The mixture was heated at 120 °C in a sealed tube for 3 hours. The mixture was diluted with 10mL water and the resultant precipitate was collected by filtration, washed with water, and recrystallized from chloroform/benzene to provide 5-bromo-1-methyl-3-(4-piperazin-1-yl-benzenesulfonyl)-1H-indole (0.302 g, 0.695mmol), M+H 436.

Similarly following the procedure described above in Example 3, but replacing 5-bromindole in Step 1 with other appropriate indoles, the following compounds were prepared:

- 1-Methyl-3-(4-piperazin-1-yl-benzenesulfonyl)-1H-indole; and;
- 6-Chloro-1-methyl-3-(4-piperazin-1-yl-benzenesulfonyl)-1H-indole.

Similarly following the procedure described above in Example 3, but skipping Step 3 and proceeding directly to Step 4 (and using various appropriate indoles in Step 1), the following compounds were prepared:

- 5-Bromo-3-(4-piperazin-1-yl-benzenesulfonyl)-1H-indole;
- 3-(4-Piperazin-1-yl-benzenesulfonyl)-1H-indole;
- 6-Chloro-3-(4-piperazin-1-yl-benzenesulfonyl)-1H-indole;
- 4-Chloro-3-(4-piperazin-1-yl-benzenesulfonyl)-1H-indole;
- 5-Methoxy-3-(4-piperazin-1-yl-benzenesulfonyl)-1H-indole;
- 5-Chloro-3-(4-piperazin-1-yl-benzenesulfonyl)-1H-indole;
- 7-Chloro-3-(4-piperazin-1-yl-benzenesulfonyl)-1H-indole;
- 7-Bromo-3-(4-piperazin-1-yl-benzenesulfonyl)-1H-indole;
- 4-Bromo-3-(4-piperazin-1-yl-benzenesulfonyl)-1H-indole;
- 5-Bromo-3-(3-chloro-4-piperazin-1-yl-benzenesulfonyl)-1H-indole;
- 5-Iodo-3-(4-piperazin-1-yl-benzenesulfonyl)-1H-indole; and
- 6-Bromo-3-(4-piperazin-1-yl-benzenesulfonyl)-1H-indole.

Similarly following the procedure described above in Example 3, but skipping Step 3 and proceeding directly to Step 4, and replacing piperazine with N-methylpiperazine (using various appropriate indoles in Step 1), the following compounds were prepared:

- 3-[4-(4-methyl-piperazin-1-yl)-benzenesulfonyl]-1*H*-indole;  
7-Bromo-3-[4-(4-methyl-piperazin-1-yl)-benzenesulfonyl]-1*H*-indole;  
6-Bromo-3-[4-(4-methyl-piperazin-1-yl)-benzenesulfonyl]-1*H*-indole;  
5-Iodo-3-[4-(4-methyl-piperazin-1-yl)-benzenesulfonyl]-1*H*-indole; and  
5 4-Bromo-3-[4-(4-methyl-piperazin-1-yl)-benzenesulfonyl]-1*H*-indole.

Similarly following the procedure described above in Example 3, skipping Step 3 and proceeding directly to Step 4, using various appropriate indoles in Step 1, and replacing 4-fluorophenyl disulfide in Step 1 with the appropriate disulfides, the following compounds were prepared:

- 10 5-Chloro-3-(3-chloro-4-piperazin-1-yl-benzenesulfonyl)-1*H*-indole;  
5-Chloro-3-(2-chloro-4-piperazin-1-yl-benzenesulfonyl)-1*H*-indole;  
3-(2-chloro-4-piperazin-1-yl-benzenesulfonyl)-1*H*-indole;  
3-(3-chloro-4-piperazin-1-yl-benzenesulfonyl)-1*H*-indole; and  
5-Bromo-3-(2-chloro-4-piperazin-1-yl-benzenesulfonyl)-1*H*-indole.

- 15 Similarly following the procedure described above in Example 3, skipping Step 3 and proceeding directly to Step 4 (and replacing piperazine with N-methylpiperazine), using various appropriate indoles in Step 1, and replacing 4-fluorophenyl disulfide in Step 1 with the appropriate phenyl disulfides, the following compounds were prepared:

- 20 5-Bromo-3-[2-chloro-4-(4-methyl-piperazin-1-yl)-benzenesulfonyl]-1*H*-indole;  
5-Chloro-3-(3-chloro-4-(4-methyl-piperazin-1-yl)-benzenesulfonyl)-1*H*-indole;  
5-Bromo-3-(3-chloro-4-(4-methyl-piperazin-1-yl)-benzenesulfonyl)-1*H*-indole;  
3-(3-chloro-4-(4-methyl-piperazin-1-yl)-benzenesulfonyl)-1*H*-indole;  
5-Chloro-3-(2-chloro-4-(4-methyl-piperazin-1-yl)-benzenesulfonyl)-1*H*-indole;  
3-(2-chloro-4-(4-methyl-piperazin-1-yl)-benzenesulfonyl)-1*H*-indole; and  
25 5-Bromo-3-(2-chloro-4-(4-methyl-piperazin-1-yl)-benzenesulfonyl)-1*H*-indole.

#### Example 4

Pharmaceutical preparations for delivery by various routes are formulated as shown in the following Tables. "Active ingredient" or "Active compound" as used in the Tables means one or more of the Compounds of Formula I.

Composition for Oral Administration

Ingredient	% wt./wt.
Active ingredient	20.0%
Lactose	79.5%
Magnesium stearate	0.5%

The ingredients are mixed and dispensed into capsules containing about 100 mg each; one capsule would approximate a total daily dosage.

5

Composition for Oral Administration

Ingredient	% wt./wt.
Active ingredient	20.0%
Magnesium stearate	0.5%
Crosscarmellose sodium	2.0%
Lactose	76.5%
PVP (polyvinylpyrrolidone)	1.0%

The ingredients are combined and granulated using a solvent such as methanol. The formulation is then dried and formed into tablets (containing about 20 mg of active compound) with an appropriate tablet machine.

10

Composition for Oral Administration

Ingredient	<u>Amount</u>
Active compound	1.0 g



- 40 -

Fumaric acid	0.5 g
Sodium chloride	2.0 g
Methyl paraben	0.15 g
Propyl paraben	0.05 g
Granulated sugar	25.5 g
Sorbitol (70% solution)	12.85 g
Veegum K (Vanderbilt Co.)	1.0 g
Flavoring	0.035 ml
Colorings	0.5 mg
Distilled water	q.s. to 100 ml

The ingredients are mixed to form a suspension for oral administration.

#### Parenteral Formulation

Ingredient	% wt./wt.
Active ingredient	0.25 g
Sodium Chloride	qs to make isotonic
Water for injection	100 ml

- 5        The active ingredient is dissolved in a portion of the water for injection. A sufficient quantity of sodium chloride is then added with stirring to make the solution isotonic. The solution is made up to weight with the remainder of the water for injection, filtered through a 0.2 micron membrane filter and packaged under sterile conditions.

Suppository Formulation

Ingredient	% wt./wt.
Active ingredient	1.0%
Polyethylene glycol 1000	74.5%
Polyethylene glycol 4000	24.5%

The ingredients are melted together and mixed on a steam bath, and poured into molds containing 2.5 g total weight.

5

Topical Formulation

Ingredients	grams
Active compound	0.2-2
Span 60	2
Tween 60	2
Mineral oil	5
Petrolatum	10
Methyl paraben	0.15
Propyl paraben	0.05
BHA (butylated hydroxy anisole)	0.01
Water	q.s. 100

All of the ingredients, except water, are combined and heated to about 60°C with stirring. A sufficient quantity of water at about 60°C is then added with vigorous stirring to emulsify the ingredients, and water then added q.s. about 100 g.

- Several aqueous suspensions containing from about 0.025-0.5 percent active compound are prepared as nasal spray formulations. The formulations optionally contain inactive ingredients such as, for example, microcrystalline cellulose, sodium carboxymethylcellulose, dextrose, and the like. Hydrochloric acid may be added to adjust pH. The nasal spray formulations may be delivered via a nasal spray metered pump typically delivering about 50-100 microliters of formulation per actuation. A typical dosing schedule is 2-4 sprays every 4-12 hours.

### Example 5

#### Radioligand binding studies

- The binding activity of compounds of this invention *in vitro* was determined as follows.

Duplicate determinations of ligand affinity are made by competing for binding of [3H]LSD in cell membranes derived from HEK293 cells stably expressing recombinant human 5-HT6 receptor.

- All determinations are made in assay buffer containing 50 mM Tris-HCl, 10mM MgSO<sub>4</sub>, 0.5 mM EDTA, 1 mM ascorbic acid, pH 7.4 at 37 °C, in a 250 microliter reaction volume. Assay tubes containing [3H] LSD (5nM), competing ligand, and membrane are incubated in a shaking water bath for 60 min. at 37°C, filtered onto Packard GF-B plates (pre-soaked with 0.3% PEI) using a Packard 96 well cell harvester and washed 3 times in ice cold 50 mM Tris-HCl. Bound [3H] LSD is determined as radioactive counts per minute using Packard TopCount.

Displacement of [3H]LSD from the binding sites was quantified by fitting concentration-binding data to a 4-parameter logistic equation:

$$\text{binding} = \text{basal} + \left( \frac{\text{Bmax} - \text{basal}}{1 + 10^{-\text{Hill}(\log[\text{ligand}] - \log \text{IC}_{50})}} \right)$$

- where Hill is the Hill slope, [ligand] is the concentration of competing radioligand and IC<sub>50</sub> is the concentration of radioligand producing half-maximal specific binding of radioligand. The specific binding window is the difference between the Bmax and the basal parameters.


- Using the procedure of this Example, Compounds of Formula I were tested and found to be selective 5-HT6 antagonists as shown below:

#	STRUCTURE	NAME	pKi	Example
1		5-Fluoro-1-(4-piperazin-1-yl-benzenesulfonyl)-1H-indole	9.0	1
4		1-(4-Piperazin-1-yl-benzenesulfonyl)-1H-indole	9.05	1
21		5-Iodo-3-[4-(4-methyl-piperazin-1-yl)-benzenesulfonyl]-1H-indole	9.15	3
26		3-(2-Chloro-4-piperazin-1-yl-benzenesulfonyl)-1H-indole	9.32	3
27		3-(3-Chloro-4-piperazin-1-yl-benzenesulfonyl)-1H-indole	9.36	3
29		5-Bromo-3-(2-chloro-4-piperazin-1-yl-benzenesulfonyl)-1H-indole	9.54	3
35		5-Chloro-3-(2-chloro-4-piperazin-1-yl-benzenesulfonyl)-1H-indole	9.61	3

While the present invention has been described with reference to the specific embodiments thereof, it should be understood by those skilled in the art that various changes may be made and equivalents may be substituted without departing from the true spirit and scope of the invention. In addition, many modifications may be made to adapt a particular situation, material, composition of matter, process, process step or steps, to the objective spirit and scope of the present invention. All such modifications are intended to be within the scope of the claims appended hereto.


Claims

1. A compound of the formula:




wherein:

- 5      $R^1$  and  $R^2$  each independently is hydrogen, alkyl, aryl, acyl, halo, nitro, amino, cyano, alkoxy, hydroxy, aryloxy, alkylthio, arylthio, thiol, carbonylamino, aminocarbonyl, or haloalkyl;
- $R^3$  and  $R^4$  each independently is hydrogen, halo, alkyl, acyl, aryl, or arylalkyl;
- $n$  is 0 to 4;
- 10     $R^6$  in each independent occurrence is hydrogen, alkyl, alkoxy, or halo;
- $R^7$  and  $R^8$  each independently is hydrogen or alkyl; and
- $R^9$  is hydrogen, alkyl, or arylalkyl;
- or an individual isomer, a racemic or non racemic mixture of isomers, a prodrug, or a pharmaceutically acceptable salt or solvate thereof.
- 15    2. The compound of formula I of claim 1, wherein:
- $R^1$  and  $R^2$  each independently is hydrogen, halo, or alkoxy;
- $R^3$  and  $R^4$  each independently is hydrogen, halo or alkyl;
- $n$  is 1 and  $R^6$  is hydrogen or halo;
- $R^7$  and  $R^8$  are hydrogen; and
- 20     $R^9$  is hydrogen or alkyl.
3. The compound of formula I of claim 1, wherein said compound has the formula:




wherein  $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^4$ ,  $n$ ,  $R^6$ ,  $R^7$ ,  $R^8$ , and  $R^9$  are defined in claim 1.

4. The compound of formula I of claim 1, wherein said compound has the formula:



wherein  $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^4$ ,  $n$ ,  $R^6$ ,  $R^7$ ,  $R^8$ , and  $R^9$  are defined in claim 1.

5. The compound of formula I of claim 1, wherein said compound has the formula:



wherein  $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^4$ ,  $n$ ,  $R^6$ ,  $R^7$ ,  $R^8$ , and  $R^9$  are defined in claim 1.

6. The compound of formula Ia of claim 3, wherein  $R^1$  and  $R^2$  each independently is hydrogen, halo or alkoxy,  $R^3$  is hydrogen or halo;  $R^4$  is hydrogen or alkyl;  $R^6$ ,  $R^7$  and  $R^8$  are hydrogen and  $R^9$  is hydrogen or alkyl.

7. The compound of formula Ia of claim 6, wherein  $R^1$  and  $R^2$  each independently is hydrogen, fluoro, bromo or methoxy;  $R^3$  is hydrogen or chloro;  $R^4$  each is hydrogen or methyl;  $R^6$ ,  $R^7$  and  $R^8$  are hydrogen and  $R^9$  is hydrogen or methyl.

8. The compound of formula Ia of claim 7, which compound is:

5-Fluoro-1-(4-piperazin-1-yl-benzenesulfonyl)-1*H*-indole;  
4-bromo-1-(4-piperazin-1-yl-benzenesulfonyl)-1*H*-indole;  
5-methoxy-1-(4-piperazin-1-yl-benzenesulfonyl)-1*H*-indole;  
1-(4-piperazin-1-yl-benzenesulfonyl)-1*H*-indole;  
5-fluoro-1-[4-(4-methyl-piperazin-1-yl)-benzenesulfonyl]-1*H*-indole;  
3-chloro-5-methoxy-2-methyl-1-(4-piperazin-1-yl-benzenesulfonyl)-1*H*-indole.

9. The compound of formula Ib of claim 4, wherein  $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^4$ ,  $R^6$ ,  $R^7$ ,  $R^8$  and  $R^9$  are hydrogen.

10. The compound of formula Ib of claim 9, which compound is:

2-(4-Piperazin-1-yl-benzenesulfonyl)-1*H*-indole.


11. The compound of formula Ic of claim 5, wherein  $R^1$  and  $R^2$  each independently is hydrogen, halo or alkoxy;  $R^3$  is hydrogen;  $R^4$  is hydrogen or alkyl;  $n$  is 1 and  $R^6$  is hydrogen or halo;  $R^7$  and  $R^8$  are hydrogen and  $R^9$  is hydrogen or alkyl.

12. The compound of formula Ic of claim 11, wherein  $R^1$  and  $R^2$  each independently is hydrogen, fluoro, chloro, bromo, iodo or methoxy;  $R^3$  is hydrogen;  $R^4$  is hydrogen or methyl;  $n$  is 1 and  $R^6$  is hydrogen or chloro;  $R^7$  and  $R^8$  are hydrogen and  $R^9$  is hydrogen or methyl.

13. The compound of formula Ic of claim 12, which compound is:


5-Bromo-1-methyl-3-(4-piperazin-1-yl-benzenesulfonyl)-1*H*-indole;  
5-bromo-3-(4-piperazin-1-yl-benzenesulfonyl)-1*H*-indole;  
3-(4-piperazin-1-yl-benzenesulfonyl)-1*H*-indole;  
1-methyl-3-(4-piperazin-1-yl-benzenesulfonyl)-1*H*-indole;  
6-chloro-3-(4-piperazin-1-yl-benzenesulfonyl)-1*H*-indole;  
4-chloro-3-(4-piperazin-1-yl-benzenesulfonyl)-1*H*-indole;  
5-methoxy-3-(4-piperazin-1-yl-benzenesulfonyl)-1*H*-indole;  
5-chloro-3-(4-piperazin-1-yl-benzenesulfonyl)-1*H*-indole;  
7-chloro-3-(4-piperazin-1-yl-benzenesulfonyl)-1*H*-indole;  
6-chloro-1-methyl-3-(4-piperazin-1-yl-benzenesulfonyl)-1*H*-indole;  
3-[4-(4-methyl-piperazin-1-yl)-benzenesulfonyl]-1*H*-indole;  
7-bromo-3-[4-(4-methyl-piperazin-1-yl)-benzenesulfonyl]-1*H*-indole;  
7-bromo-3-(4-piperazin-1-yl-benzenesulfonyl)-1*H*-indole;  
7-bromo-3-(4-piperazin-1-yl-benzenesulfonyl)-1*H*-indole;

- 6-bromo-3-[4-(4-methyl-piperazin-1-yl)-benzenesulfonyl]-1H-indole;  
 5-iodo-3-[4-(4-methyl-piperazin-1-yl)-benzenesulfonyl]-1H-indole;  
 4-bromo-3-[4-(4-methyl-piperazin-1-yl)-benzenesulfonyl]-1H-indole;  
 4-bromo-3-(4-piperazin-1-yl-benzenesulfonyl)-1H-indole;  
 5-iodo-3-(4-piperazin-1-yl-benzenesulfonyl)-1H-indole;  
 6-bromo-3-(4-piperazin-1-yl-benzenesulfonyl)-1H-indole;  
 3-(2-chloro-4-piperazin-1-yl-benzenesulfonyl)-1H-indole;  
 3-(3-chloro-4-piperazin-1-yl-benzenesulfonyl)-1H-indole;  
 5-bromo-3-[2-chloro-4-(4-methyl-piperazin-1-yl)-benzenesulfonyl]-1H-indole;  
 5-bromo-3-(2-chloro-4-piperazin-1-yl-benzenesulfonyl)-1H-indole;  
 5-chloro-3-(3-chloro-4-piperazin-1-yl-benzenesulfonyl)-1H-indole;  
 5-chloro-3-(3-chloro-4-(4-methyl-piperazin-1-yl)-benzenesulfonyl)-1H-indole;  
 3-(3-chloro-4-(4-methyl-piperazin-1-yl)-benzenesulfonyl)-1H-indole;  
 5-bromo-3-(3-chloro-4-(4-methyl-piperazin-1-yl)-benzenesulfonyl)-1H-indole;  
 5-bromo-3-(3-chloro-4-piperazin-1-yl-benzenesulfonyl)-1H-indole;  
 5-chloro-3-(2-chloro-4-piperazin-1-yl-benzenesulfonyl)-1H-indole;  
 5-chloro-3-(2-chloro-4-(4-methyl-piperazin-1-yl)-benzenesulfonyl)-1H-indole;  
 3-(2-chloro-4-(4-methyl-piperazin-1-yl)-benzenesulfonyl)-1H-indole;  
 5-bromo-3-(2-chloro-4-(4-methyl-piperazin-1-yl)-benzenesulfonyl)-1H-indole.
14. A process for producing a compound of formula I of claim 1, comprising contacting a 4-halobenzenesulfonyl-indole of the formula:



wherein X is halo,


with a piperazine of the formula:



to produce a compound of the formula:




- 48 -




wherein  $n$ ,  $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^4$ ,  $R^6$ ,  $R^7$ ,  $R^8$  and  $R^9$  are as defined in claim 1.

15. A process for producing a compound of formula Ia of claim 3, comprising contacting formula




wherein X is halo,

with a piperazine of the formula:




to produce a compound of the formula:




wherein  $n$ ,  $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^4$ ,  $R^6$ ,  $R^7$ ,  $R^8$  and  $R^9$  are as defined in claim 1.

16. A process for producing a compound of formula Ib of claim 4, comprising contacting formula




wherein X is halo,

5 with a piperazine of the formula:




to produce a compound of the formula:



wherein n, R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup>, R<sup>6</sup>, R<sup>7</sup>, R<sup>8</sup> and R<sup>9</sup> are as defined in claim 1.


10 17. A process for producing a compound of formula Ic of claim 5, comprising contacting formula




wherein X is halo,

with a piperazine of the formula:

- 50 -



to produce a compound of the formula:



wherein  $n$ ,  $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^4$ ,  $R^6$ ,  $R^7$ ,  $R^8$  and  $R^9$  are as defined in claim 1.

18. A pharmaceutical composition comprising a therapeutically effective amount of at least one compound of claim 1 to 13 in admixture with one or more pharmaceutically acceptable carrier for the treatment of diseases.
19. Use of one or more compounds of formula I of claim 1 for the manufacture of a medicament for the treatment or prevention of a disease state that is alleviated by 5-HT<sub>6</sub> agonist.
20. The use of claim 19, wherein the disease state comprises disorders of CNS.
21. The use of claim 20, wherein the disease state comprises psychoses, schizophrenia, manic depressions, neurological disorders, memory disorders, attention deficit disorder, Parkinson's disease, amyotrophic lateral sclerosis, Alzheimer's disease and Huntington's disease.
22. The use of claim 19, wherein the disease state comprises disorders of the gastrointestinal tract.
23. The use of claim 19, wherein the disease state comprises obesity.
24. The invention is hereinbefore described.

## INTERNATIONAL SEARCH REPORT

 International Application No.  
 PCT/EP 03/11323

 A. CLASSIFICATION OF SUBJECT MATTER  
 IPC 7 A61K31/404 C07D209/08 C07D209/30 A61P25/00

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

CHEM ABS Data, EPO-Internal, WPI Data, PAJ

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
E	WO 03 104193 A (HOFFMANN LA ROCHE) 18 December 2003 (2003-12-18) claims 1,21 ---	1-24
A	WO 02 41889 A (BROMIDGE STEVEN MARK ;SMITHKLINE BEECHAM PLC (GB)) 30 May 2002 (2002-05-30) see claim 1 and description, page 1, lines 19-21 ---	1-24
A	WO 02 32863 A (BIOVITRUM AB ;JOHANSSON GARY (SE); CALDIROLA PATRIZIA (SE); NILSSO) 25 April 2002 (2002-04-25) claims 1,23 --- -/-	1-24

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

\* Special categories of cited documents:

\*A\* document defining the general state of the art which is not considered to be of particular relevance

\*E\* earlier document but published on or after the international filing date

\*L\* document which may throw doubt on priority claims) or which is cited to establish the publication date of another claim or other special reason (see specified)

\*C\* document referring to an oral disclosure, use, exhibition or other means

\*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

\*S\* document member of the same patent family

Date of the actual completion of the international search

30 January 2004

Date of mailing of the international search report

11/02/2004

 Name and mailing address of the ISA  
 European Patent Office, P.B. 5818 Patentkan 2  
 NL - 2200 HV Rijswijk  
 Tel: (+31-70) 340-2040, Tx. 31 651 epo nl,  
 Fax: (+31-70) 340-3016

Authorized officer

Bérillon, L

## INTERNATIONAL SEARCH REPORT

International Application No.

PCT/EP 03/11323

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	LEE M ET AL: "5-HT <sub>6</sub> SEROTONIN RECEPTOR BINDING AFFINITIES OF N1-BENZENESULFONYL AND RELATED TRYPTAMINES" MEDICINAL CHEMISTRY RESEARCH, BIRKHAUSER, BOSTON, US, vol. 10, no. 4, 2000, pages 230-242, XP001079522 ISSN: 1054-2523 the whole document ---	1-24
A	TSAI Y ET AL: "N1-(Benzenesulfonyl)tryptamines as novel 5-HT <sub>6</sub> antagonists" BIOORGANIC & MEDICINAL CHEMISTRY LETTERS, OXFORD, GB, vol. 10, no. 20, 16 October 2000 (2000-10-16), pages 2295-2299, XP004224204 ISSN: 0960-894X the whole document -----	1-24

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/EP 03/11323

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
  
2. ☒ Claims Nos.: 1-24 (part)  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:  
see FURTHER INFORMATION sheet PCT/ISA/210
  
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
  
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
  
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
  
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

### Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

## FURTHER INFORMATION CONTINUED FROM PCT/SA/ 210

Continuation of Box I.2

Claims Nos.: 1-24 (part)

Present claims 1-24 relate to compounds defined by reference to a desirable characteristic or property, namely prodrugs. The claims cover all compounds having this characteristic or property, whereas the application provides support within the meaning of Article 6 PCT and/or disclosure within the meaning of Article 5 PCT for only a very limited number of such compounds. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible. Independent of the above reasoning, the claims also lack clarity (Article 6 PCT). An attempt is made to define the compounds by reference to a result to be achieved. Again, this lack of clarity in the present case is such as to render a meaningful search over the whole of the claimed scope impossible. Consequently, the search has been carried out for those parts of the claims which appear to be clear, supported and disclosed, namely those parts of the claims 1-24 relating to compounds of formula (I), individual isomer, racemic or non racemic mixture of isomers or pharmaceutically acceptable salt or solvate thereof.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No.

PCT/EP 03/11323

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 03104193	A	18-12-2003	WO 03104193 A1	18-12-2003
			US 2003229069 A1	11-12-2003
WO 0241889	A	30-05-2002	AU 2071502 A	03-06-2002
			WO 0241889 A2	30-05-2002
			EP 1335722 A2	20-08-2003
WO 0232863	A	25-04-2002	AU 9619301 A	29-04-2002
			BR 0114552 A	01-07-2003
			CA 2422717 A1	25-04-2002
			EP 1326830 A1	16-07-2003
			NO 20031589 A	28-05-2003
			WO 0232863 A1	25-04-2002
			US 2002165251 A1	07-11-2002